



## Certificate of Analysis

Wild Honey NZ (2017) Ltd  
 52 Woods Rd South, Waitete Bay  
 Coromandel  
 Attention: Geoff Gillott  
 Phone: 021 069 3378  
 Email: gilloty@hotmail.com

Lab Reference: 22-13242  
 Submitted by: N/A  
 Date Received: 07/04/2022  
 Testing Initiated: 7/04/2022  
 Date Completed: 7/04/2022  
 Order Number: N/A  
 Reference: N/A

### Report Comments

Samples were collected by yourselves (or your agent) and analysed as received at Analytica Laboratories. Samples were in acceptable condition unless otherwise noted on this report.  
 Specific testing dates are available on request.

### Results Summary

#### 3in1 in Honey

Laboratory ID	Sample ID	Dihydroxyacetone (DHA)	Methylglyoxal (MG/MGO)	Non-Peroxide Activity* (NPA)	Hydroxymethylfurfural (HMF)
	<i>Units Reporting Limit</i>	mg/kg 40	mg/kg 8	%w/v phenol eq. 1.3	mg/kg 1
22-13242-1	WH2110	881	538	15.4	28.6

#### 3in1 in Honey Approver:

Gurmeet Singh, Dip. Tech. (Sci)  
 Laboratory Technician

### Method Summary

#### 3in1

Determination of Dihydroxyacetone (DHA), Methylglyoxal (MG/MGO) and Hydroxymethylfurfural (HMF) by aqueous extraction, derivatisation, and UPLC (diode array) analysis in accordance with in-house procedures.

#### NPA

Non-Peroxide Activity (NPA) values are not directly measured by the laboratory, but are calculated from the measured methylglyoxal concentration in the honey according to the requirements of the client. The calculation is based on published data<sup>(†)</sup> comparing the NPA and methylglyoxal concentration measured in a range of honey samples. These calculated values are not accredited by IANZ and do not imply that the honey is or is not manuka honey. NPA values less than 5 are an estimate based on extrapolation of the relationship between methylglyoxal and NPA

*(†) Isolation by HPLC and characterisation of the bioactive fraction of New Zealand manuka (Leptospermum scoparium) honey. C. J. Adams, et al. Carbohydrate Research 343 (2008) 651-659. And, Corrigendum to "Isolation by HPLC and characterization of the bioactive fraction of New Zealand manuka (Leptospermum scoparium) honey" [Carbohydr. Res. 343 (2008) 651]. Carbohydrate Research 344 (2009) 2609. C. J. Adams, et al.*